

Re-Construction of Paleo-Sedimentation Processes of Aquifers Underlying Igueben Using Geo-Electrical Resistivity Signature and Borehole Data

Samuel O. Salufu¹, Isaac Aigbedion^{1*}

¹Geophysics/Physics Department, Ambrose Alli University, Ekpoma, Nigeria

INFORMATION

Article history

Received 18 July 2021

Revised 21 December 2021

Accepted 22 December 2021

Keywords

Aquifer

Palaeo-sedimentation processes

Geo-electrical resistivity

Palaeo-Sedimentation Processes

Igueben

Contact

*Isaac Aigbedion

isaacaigbedion@yahoo.com

ABSTRACT

Igueben is located in the central of Edo State, Nigeria. It has neighboring towns; Ogwa, Ugbegun, Ugiogba, Ebelle, and Ekpon. These towns have the same aquifer but significant variations in the water table. However, this study is intended to unravel the salient ancient hydrological processes that occurred in the study area which resulted in this variation. Vertical electrical sounding (VES) data and borehole data of the area are integrated to generate geo-electrical resistivity section of the area, to identify the aquifer trends, body, and shape to re-construct the paleo-sedimentation processes of the aquifers that underlie Iguebe and the neighboring towns. The inversion model result for the seven VES conducted in the study area show that the curve types are HQA, AHA, A, AHA, AHA, HAQ, and AQA, respectively. These curve type suggested that the aquifer type is an arenite sandstone aquifer that has been transported from a long distance. The geo-electrical pattern and shape show that Iguebe and Ogwa (shallow marine) are the points of deposition from where other nearby towns (Ugbegun, Ugiogba, Ebelle, and Ekpon) sourced their aquifer by gravity settling from suspended sediments in water body into adjacent deep marine environment (Ugbegun, Ugiogba, Ebelle, and Ekpon). The two towns are the hydraulic-head of the aquifer in the study area. These geological processes are responsible for the variation in the water table in the study area.

1. Introduction

Paleo-sedimentation processes of an aquifer are all the processes that had taken place in the past as at the time of transportation and deposition of an aquifer in a basin (Arua 1986; Arua and Okoro, 1989). These processes are controlled by sediment source availability, agents of weathering, transport media (glacier, wind, and water), proximity of source to basin, bio-chemical activities (Stow et al., 2001; VerStraeten et al., 2011), mechanical activities, and change in sea level (Brett et al., 2011; Ryan et al., 2015). Paleo-environment where these processes occurred determines the yield of an aquifer. These processes determine the character of an aquifer (Woodrow, 1985; Gary, 2009). Hence the knowledge about the paleo-sedimentation of an aquifer is a clue to the aquifer characteristics. However, aquifer formed by these processes can be altered by geological processes such as erosion and tectonic activities. Integration of geochemical and petrographical data of sandstone in a sedimentary basin

can reveal the nature, Palaeo-environment of the sediments that fill the basins (Valloni and Mezzardi, 1984; McLennan et al., 1993; Kroonenberg, 1994).

However, Aigbedion and Salufu (2021) have been able to combine geoelectrical section of sediments and petrographic of sandstone of a basin to reconstruct the Paleo transportation and depositional environment of the sandstone with a successful result. Existing literatures has given the evolution of Anambra Basin, its stratigraphic setting, and Paleo environment using petrographic and sedimentological studies (Nwajide and Huque, 1985; Idowu and Ekweozor, 1993).

Little/scanty literatures about the hydrogeology and geology of Igueben exist. However, borehole data of the wells in the area was provided by Rock Well Drilling Company as acquired in 2018 (Rock Well Drilling Company, 2018).



Igueben has been known to have groundwater occurrence at both shallow depths (120-140 m) and deep depths (220 m and above) while Ugiogba, Ugbegun, Ebelle, and Ekpon that are neighboring towns have deep aquifers. However, no existing literature have been able to explain the reason for such occurrence. Borehole data from these towns have shown that the aquifer comprises the same derived properties such as fine grained sandstone facies. Existing literatures have not been able to give reason(s) for the variation in the water table

among the Oligocene age aquifers that exist in Igueben. This present study is intended at unraveling the reason and the salient hydrological processes that occur in the study area by re-constructing the Paleo-sedimentation processes of the aquifers that underlie Iguebe and the neighboring towns using geo-electrical section according to Koefoed (1979) and borehole data of the areas to characterize the aquifers according to aquifer's sandstone type (Aigbedion and Salufu, 2021), aquifer deposition environment and hydraulic head.




Fig. 1. Satellite map of the study area showing Igueben and neighbouring towns

2. Geographical and Local Geology Setting of Study Area

The study area is located in the part of Esan Central in Edo State. The area is part of the Lower Benue Trough (Anambra Basin) arm that extends to Edo State (Benkheilil, 1989). The basin was evolved after the Santonian Tectonic event that occurred in the Southern Nigerian (Benkheilil, 1989).

It comprises Iguebe, Ugiogba, Ugbegun, Ogwa, Ebelle, and Ekpon (Fig. 1) in Iguebe Local Government Area of Edo State, Nigeria. The area is accessible by major road, Uromi-Agbor Road, and other minor roads (Fig. 1).

The area is underlain by clayey sandstone and lateritic sandstone facies that belong to Ogwashi-Asaba Formation.

The formation has a general strike of $89^{\circ}\text{NE}-269^{\circ}\text{SW}$, dip direction is towards 178°SE , and dip is 5° . This shows that the basin is tectonically stable, no history of tectonic activities. The lateritic sandstone facies covers Ugbegun, Ugiogba, Ogwa, Igueben, and Ebelle while Ekpon is covered by clayey sandstone facies (Fig. 2). The sandstones have general dip direction of south east direction with 5° dip (Fig. 2). The sandstones are reddish to brown and friable.

3. Material and Methods

3.1. Data acquisition

Seven VES were conducted in Igueben and neighboring towns (Fig. 3) using Schlumberger array (Fig. 4) to determine the depth of water table and depth of aquifer occurrence in

each town. Borehole data of four wells in the study area were obtained from [Rock Well Drilling Company \(2018\)](#). The result of seven VES obtained in the area was integrated with the borehole data to generate geo-electrical resistivity section of the area. The motive of the geo-electrical section of the

study area was used to characterize the aquifer and infer the paleo-sedimentation processes that had taken place in the course of aquifer deposition in the area. Thus salient information useful to the modeling of the hydrogeological setting of the area was deduced.




Fig. 2. Geological map of the study area

As the fluid flow into the basin or from one part of the basin to another part in the time past, the fluid carried sediments along and got deposited into the basin as fluid mass. The fluid masses either gained or lose velocity by moving into an area where the velocity had changed in time at any position. The deposited sediments can be aquifer, aquitard or aquiclude. The total fluid acceleration during sedimentation in a basin is given as $\frac{Du}{Dt}$.

where, Capital D is total differential coordinate and s is natural coordinate.

Hence,

$$\frac{Du}{Dt} = \frac{\partial u}{\partial t} + \left(\frac{\partial u}{\partial s}\right) \quad (1)$$

Expanding into 3D coordinate [Equation 2](#) becomes:

$$\frac{Du}{Dt} = \frac{\partial u}{\partial t} + (u\frac{\partial u}{\partial x} + v\frac{\partial u}{\partial y} + w\frac{\partial u}{\partial z}) \quad (2)$$

Thus, [Equation 2](#) can be transformed to:

$$\frac{Du}{Dt} = \frac{\partial u}{\partial t} + u \cdot \nabla u \quad (3)$$

Aquifer that is deposited in a basin can be located by VES using the basic principle of ohm's law:

$$R = \frac{V}{I} \quad (4)$$

Resistivity in Equation 4 is the measured resistivity by Terrameter. The true resistivity of the subsurface is gotten by

multiplying Equation 4 with geoelectric factor (K) to have apparent resistivity (ρ_a).




Fig. 3. Topography map of the study area showing the VES point locations




Fig. 4. Schlumberger array (after Lowrie, 1997)

Table 1. Raw field VES data acquired in the study area

S/N	AB/2	MN/2	Ugbegun (Ωm)	Ugiogba (Ωm)	Ogwa (Ωm)	Igueben North (Ωm)	Iguebe South (Ωm)	Ebelle (Ωm)	Ekpon (Ωm)
1	5	2	1493	726	1493	1593	786	616.14	770
2	15	2	1559	887	1359	1659	987	826.9	1571
3	20	2	2117	1174	2217	2217	1274	918.6	2134
4	40	5	2178	1520	2378	2678	1620	1027	3145
5	80	5	3156	1493	3456	3456	1593	2091	4156
6	120	10	2267	1559	4767	4567	1659	3189	6473
7	140	10	2267	2117	4867	4667	2217	3563	7895
8	180	10	2589	2478	4989	4789	2678	3465	9312
9	220	15	4498	4356	4898	4798	3456	3387	13459
10	360	15	4821	4567	4701	4801	4567	3667	15678

$$\rho_a = K \frac{V}{I} \tag{5}$$

K is computed using the general Equation 6.

Let the separations of current and potential electrodes in Fig. 4b L and a, respectively. General equation is given as:

$$\rho_a = 2\pi \frac{V}{I} \left[\frac{1}{\left(\frac{1}{A} - \frac{1}{B}\right) - \left(\frac{1}{C} - \frac{1}{D}\right)} \right] \tag{6}$$

Then,

$$A = D = \frac{(L-2)}{2} \tag{7}$$

$$C = B = \frac{(L-2)}{2} \tag{8}$$

Substituting in Equation 6.

$$\rho_a = 2\pi \frac{V}{I} \left[\frac{1}{\left(\frac{2}{L-a} - \frac{2}{L+a}\right) - \left(\frac{2}{L+a} - \frac{2}{L-a}\right)} \right] \tag{9}$$

Hence, Equation 9 becomes:

$$\rho_a = \frac{\pi V}{4 I} \left(\frac{L^2 - a^2}{a} \right) \tag{10}$$

3.2. Inversion Model

Subsequently, Res1Dinvers is used to carry out the inversion model to produce a model response that matches the measured values by using the least-squares optimization method after Lines and Treitel (1984) according to Equation 11 as shown below:

$$(J^T J + \lambda I) \Delta_{q_k} = J^T g \tag{11}$$

The purpose of the inversion subroutine was to determine the resistivity and thickness of the aquifer in the study area and to determine depth to water table and geology of the area.




Fig. 5. a) Inverted model layer for VES 1 taken at Ugbeugun, b) Inverted model layer for VES 2 taken at Ugiogba, c) Inverted model layer for VES 3 taken at Ogwa and d) Inverted model layer for VES 4 taken at Iguebe north in Edo State

Where; Q is the model parameter vector that consists of the logarithm of the resistivity and thickness of the layers, g is the

discrepancy vector that consists of the difference between the logarithms of the calculated and measured apparent

resistivity values, Δq is the model parameter change vector, and J is the Jacobian matrix of partial derivatives. The elements of Jacobian matrix are expressed in Equation 12.

$$J_{ij} = \frac{\partial f_i}{\partial q_j} \tag{12}$$

It is the change in the i^{th} model response f_i due to a change in

the j^{th} model parameter q_j . The I is the identity matrix. The factor λ is known as the Marquardt or damping factor, and this method is also known as the ridge regression method (Inman, 1975). The damping factor effectively constrains the range of values that the components of parameter change vector can Δq take. The damped least-squares method attempts to minimize a combination of the magnitude of the discrepancy vector and the parameter change vector.




Fig. 6. a) Inverted model layer for VES 5 taken at Iguebe south, b) Inverted model layer for VES 6 taken at Ebelle and c) Inverted model layer for VES 7 taken at Ekpon in Edo State

4. Results and Discussion

The result of raw field resistivity data acquired in the study area is given in Table 1. The inversion model is given in Fig. 5 and Fig. 6. The inversion model for the seven VES conducted in the study area showed that the curve types for the seven VES are HQA, AHA, A, AHA, AHA, HAQ, and AQA, respectively. The pattern of the curve type showed that the aquifers are arenitic sandstone that had undergone second cycle, sediments recycled probably from a long distance. The resistivity results for the seven VES acquired in the study area showed that Ogwa and Iguebe have the shallowest water table 110 m and 120, respectively. However, Iguebe result indicated that Iguebe has two distinct water table; shallow and deep; 120 m and 220 m, respectively. The shallow water

table is restricted to the northern part of Iguebe while the deep water table occurs in the southern part. Ugiogba, Ugbegun, Ebelle, and Ekpon, have deep water table; 240 m, 250 m, 230 m, and 240 m, respectively.

5. Re-Construction of Paleo-Sedimentation Processes of Iguebe and Environs

The integration of resistivity section and the borehole data in the study area obviously revealed the fact that the aquifer deposits depth was shallow at Iguebe north and Ogwa, thus became deepening towards the south and extreme north of the study area in Iguebe south, Ugbegun, Ugiogba, Ebelle, and Ekpon. They are located as shown in the geo-electrical resistivity section of the study area (Fig. 7).

The borehole data (Fig. 8) in the study area confirmed this fact. The pattern of the geo-electrical section showed that

Ogwa and Iguebe north represented the point in the ancient sea where river flowed into the standing body of water (sea).




Fig. 7. Geo-electrical resistivity section of the study area




Fig.8. Litho-log of boreholes section of the study area (Rock Well Drilling Company, 2018)

The geo-electric pattern suggests that Ogwa and Igueben are the hydraulic-head of the aquifer in the study area. The sediment continued to get deposited at the mouth of the standing sea and subsequently got distributed to deeper part (Ugiogba, Ugbegun, Iguebe south, Ebelle, and Ekpon) of the ancient sea by the flow of the river into the sea, in response to the gravity.

The grain size of very fine grained sandstone and siltstone sequence (Fig. 8) that majorly the character of the aquifers in the study area corroborated the fact that the sandstone were

transported from very far distance into the basin (sea) later moved from the mouth of the sea (shallow part) to other deeper parts as suspended load within the water body (sea). The motive of the geo-electrical section and litho-log of boreholes section indicates Ogwa and Iguebe north to be the hydraulic head (HH) of the entire aquifers in the study area (Fig. 8). This is due to the manner the paleo-sedimentation processes occurred in the area. Thus, groundwater moved from the Ogwa and Igueben north radially to other aquifers around them. Hence the aquifer that underlies the study area was deposited in a transitional (tidal flat) between marine and

fluvial where wave energy washed silt and clay away, leaving sandstone particles behind.

6. Conclusion

Palaeo-sedimentation processes of Iguebe and environs have been successfully carried out using integrated data of vertical electrical resistivity and borehole in the area. The data was used to generate geo-electrical resistivity section of the aquifers that underlay the area in order to re-construct the aquifers trends and body. Subsequently, the ancient geological processes that led to the deposition of the aquifers and the palaeo-environment of the aquifer was delineated using the geo-electric section and the borehole data. The study has shown that Iguebe north and Ogwa are the depositor where wave energy from marine environment washed the transported sandstone from far source and distributed it to deeper parts (Iguebe south, Ugbegun, Ugiogba, Ebelle, and Ekpon) of the marine by suspension of particles that settled down due to gravity effect. Thus, the aquifers were deposited at different marine environments during Oligocene. This was responsible for the variations in the water table across the study area.

References

- Aigbedion, I., Salufu, S., 2021. Impact of Sandstone Maturity on Groundwater Resistivity in Geophysical Groundwater Exploration in Parts of Edo Central and Agbor, Delta State, Nigeria. *South Asian Research Journal of Natural Products* 4 (2), 1-10.
- Arua, I., 1986. Palaeo-environment of Eocen Deposits in Afikpo Syncline, Southern Nigeria. *Journal of African Sciences* 5, 297-284.
- Arua, I., Okoro, A.O., 1989. Reconstruction of Palaeo-wave and paleo-depth Regimes of Nkporo Sea (Campano-Maastrichtian) Using Wave formed Ripple Marks. *Journal of Mining and Geology* 25 (1-2), 75-79.
- Benkhelil, J. 1989. The Evolution of the Cretaceous Benue Trough, Nigeria. *Journal of African Earth Sciences* 8, 251-282.
- Brett, C.E., Baird, G.C., Bartholomew, A.J., Desantis, M.K., VerStraeten, C.A., 2011. Sequence stratigraphy and a revised sea-level curve for the Middle Devonian of eastern North America. *Palaeogeography, Palaeoclimatology, Palaeoecology* 304 (1-2), 21-53.
- Gary, N., 2009. *Sedimentology and Stratigraphy* Second Edition Wiley Black publisher.
- Idowu, J.O., Ekweozor, C.M., 1993. Petroleum potential of Cretaceous shales in the Upper Benue Trough, Nigeria. *Journal of Petroleum Geology* 16, 249-264.
- Inman, J.R., 1975. Resistivity inversion with ridge regression. *Geophysics* 40, 798-817.
- Koefoed, O., 1979. *Geosounding Principles 1: Resistivity sounding measurements*. Elsevier Science Publishing Company, Amsterdam.
- Kroonenberg, S.B., 1994. Effects of provenance, sorting and weathering on the geochemistry of fluvial sands from different tectonic and climatic environment. *Proceedings of the 29 International Geological Congress, Part A*, pp. 69-81.
- Lines, L.R., Treitel, S., 1984. Tutorial: A review of least-squares inversion and its application to geophysical problems. *Geophysical Prospecting* 32, 159-186.
- Lowrie, W., 1997. *Fundamentals of Geophysics*. Institute of Geophysics, Swiss Federal Institute of Technology. Zurich, Switzerland. Cambridge University Press. 354 p.
- McLennan, S.M., Hemming, S., McDaniel, D.K., Hanson, G.N., 1993. Geochemical approaches to sedimentation, provenance and tectonics, In: Johnson, M.J. and Basu, A. (eds.): *Geological Society of America, Special Paper* 285, 21-40.
- Nwajid, C.S., Hoque M., 1985. Tectono-sedimentological evolution of an elongate intracratonic basin (aulacogen): The case of the Benue Trough of Nigeria. *Journal of Mineralogy and Geology* 21 (1-2), 19-26.
- Rock Well Drilling Company, 2018. Unpublished article, Auchi-Edo State-Nigeria.
- Ryan, D.W., Juergen, S., 2015. Sedimentary Facies and Depositional Environment of the Middle Devonian Formation of New York, USA. *Journal of Sedimentary Research* 85, 1393-1415.
- Stow, D.A.V, Huc, A.Y., Bertrand, P., 2001. Depositional processes of black shales in deep water: *Marine and Petroleum Geology* 18, 491-498.
- Valloni, R., Mezzardi, G., 1984. Compositional suites of terrigenous deep-sea sands of the present continental margins. *Sedimentology* 31, 353-364.
- VerStraeten, C.A., Brett, C.E., Sageman, B.B., 2011. Mudrock sequence stratigraphy: a multi-proxy (sedimentological, paleobiological and geochemical) approach, Devonian Appalachian Basin: *Palaeogeography, Palaeoclimatology, Palaeoecology* 304, 54-73.
- Woodrow, D.L., 1985. Paleogeography, paleoclimate, and sedimentary processes of the Late Devonian Catskill Delta, in Woodrow, D.L., and Sevon, W.D., eds., *The Catskill Delta: Geological Society of America, Special Paper* 201, p. 51-63.